



Conference Abstract

P.67 The Effects of Chemotherapy on Arterial Inflammation Assessed by ¹⁸FDG PET-CT in Patients with Lymphoma

Constantinos Anagnostopoulos², Stavroula Giannouli³, Nikolaos Ioakimidis¹, Paulos Kafouris⁴, Iosif Koutagiari¹, Anastasia Sioni⁵, Eirini Solomou^{1,*}, Dimitrios Terentes-Printzios¹, Dimitrios Tousoulis¹, Charalampos Vlachopoulos¹

¹Hippokraton General Hospital, 1st Cardiology Department, Athens Medical School

²Academy of Athens Biomedical Research Foundation, Center for Experimental Surgery, Clinical and Translational Research, Biomedical Research Foundation

³Academy of Athens Biomedical Research Foundation, Center of Systems Biology

⁴Hippokraton General Hospital, Department of Hematology

⁵Academy of Athens Biomedical Research Foundation, Center of Systems Biology

Keywords

Lymphoma
chemotherapy
arterial-inflammation
¹⁸FDG PET-CT

ABSTRACT

Introduction: Anti-cancer treatment can lead to increased cardiovascular morbidity among lymphoma survivors [1]. This may be the result of direct effect of treatment on heart function, or indirect acceleration of atherosclerosis. ¹⁸F-fluorodeoxyglucose (FDG) uptake is a sensitive and robust marker for assessment of atherosclerotic inflammation [2,3,4].

Purpose: To investigate the effects of chemotherapy on arterial inflammation using FDG-PET CT in patients with lymphoma.

Methods: Fifty nine (mean age 58 ± 17 years) patients with Hodgkin (*n* = 39) or non-Hodgkin lymphomas (*n* = 20) underwent ¹⁸FDG PET-CT imaging at baseline, interim and after completion of chemotherapy as part of their routine protocol. Arterial inflammation was assessed by arterial target to background ratio (TBR) of the aortic wall along the entire aorta. The index vessel TBR (the vessel with the higher value at baseline) was used for assessment of arterial inflammation. Patients with Hodgkin Lymphomas (HL) underwent therapy with Doxorubicin, bleomycin, vinblastine and dacarbazine (ABVD). The interim of their treatment was set at 1 to 3 days prior to initiating the 3rd chemotherapy cycle. Patients with non Hodgkin Lymphomas (NHL) underwent therapy with cyclophosphamide, doxorubicin, vincristine, and prednisone + rituximab (R-CHOP). The interim of their treatment was set at 2 weeks post the 4th chemotherapy cycle. All patients were reassessed 6 weeks after chemotherapy completion.

Results: There were no differences in age and atherosclerotic risk factors (hypertension, diabetes, dyslipidemia and smoking), between the two groups (all *p* > 0.05). Similarly, there were no differences in mean (± SD) index vessel TBR between HL and NHL patients (2.4 ± 0.7 vs 2.7 ± 0.9, respectively, *p* = 0.65). In the whole study population the index vessel TBR progressively decreased after the end of therapy (by 0.53 ± 0.11, from baseline to 6 weeks following the end of therapies) (*F* = 10.94, *p* < 0.001, ANOVA). The index vessel TBR decreased in both HL and NHL patients at 6 weeks after therapy compared to baseline level (all *p* < 0.01, ANOVA, Figure). The decrease at the interim scan was more pronounced in NHL compared to HL patients, however at 6 weeks after chemotherapy completion the index vessel TBR decreased further in patients with HL, while it increased slightly compared to interim levels in NHL patients (Figure).

Conclusion: Arterial inflammation is reduced during and post-chemotherapy in patients with lymphoma. The index vessel TBR changes at the interim phase and 6 weeks after therapy completion indicate a different effect of specific treatment regimes in arterial inflammation between HL and NHL patients.

REFERENCES

- [1] Boyne DJ, Mickle AT, Brenner DR, Friedenreich CM, Cheung WY, Tang KL, et al. Long-term risk of cardiovascular mortality in lymphoma survivors: a systematic review and meta-analysis. *Cancer Med* 2018;7:4801–13.
- [2] Rosenbaum D, Millon A, Fayad ZA. Molecular imaging in atherosclerosis: FDG PET. *Curr Atheroscler Rep* 2012;14:429–37.
- [3] McKenney-Drake ML, Moghbel MC, Paydary K, Alloosh M, Houshmand S, Moe S, et al. ¹⁸F-NaF and ¹⁸F-FDG as molecular probes in the evaluation of atherosclerosis. *Eur J Nucl Med Mol Imaging* 2018;45:2190–200.
- [4] Toutouzias K, Koutagiari I, Benetos G, Aggeli C, Georgakopoulos A, Athanasiadis E, et al. Inflamed human carotid plaques evaluated by PET/CT exhibit increased temperature: insights from an *in vivo* study. *Eur Heart J Cardiovasc Imaging* 2017;18:1236–44.

© 2020 Association for Research into Arterial Structure and Physiology. Publishing services by Atlantis Press International B.V. This is an open access article distributed under the CC BY-NC 4.0 license (<http://creativecommons.org/licenses/by-nc/4.0/>).